



Pecan rootstock reactions in a cotton root rot nursery

2013-16

TAMU Research & Extension Center Uvalde

Mark Black, Alfred Sanchez, Larry Stein, Kimberly Cochran

USDA ARS Pecan Breeding & Genetics, Somerville

L. J. Grauke and Xinwang Wang

Texas A&M University, College Station

Patricia Klein with Nolan Bentley

Funded in part by **Tex. Dept. Agric. & USDA**
Specialty Crop Block Grant

Cotton







Cotton root rot disease

- Native fungus, *Phymatotrichopsis omnivora*
- SW USA, northern Mexico
- High temperature strain of a pathogen in the last Ice Age forests?
- Essentially one clone
- Wide dicot host range
- Native dicot plants have some resistance

“As individual trees have shown resistance toward the disease, some authorities are in hopes of securing strains immune or highly resistant of the trouble.”

J.H. Burkett, 1922 TDA Bulletin

CRR risk factors

- Calcareous high pH soils
- High soil temperature in summer
- Periods of soil saturation: rains, irrigations
- Any stress that stops summer root growth
- Close plant spacing
- Monoculture with susceptible rootstock
- Heavy fruit load








Apple

Cotton root rot of Pecan





Mary Olsen, Univ. of Arizona




Dept. of Plant Pathology
University of Arizona



Spore mats on soil




Releases mating (?)
spores



Phymatotrichopsis omnivora distribution in USA and Mexico

B.S. Richter 2007 <http://www.cals.ncsu.edu/course/pp728/Phymatotrichopsis/>



Coahuila south to
Jalisco and Veracruz

Carya illinoensis native range.

http://en.wikipedia.org/wiki/File:Carya_illinoensis_map.png

Positives

- Pecan evolved in Texas and NE Mexico; challenged by CRR fungus; great diversity among natives and named varieties
- In general, pecan is considered moderately resistant
- Rootstocks already used 100% for improved varieties




Figure 3. Seed stocks preferred by pecan nurserymen within regions.


Strategy

- Seedlings from N, E, S, W and Mixed provenances
- Site history of CRR
- Close plantings (2 ft) of 10-plants/plot
- 8 replications (fungus not evenly distributed)
- Ramped up disease by planting alfalfa
- Subsurface drip irrigation





Pecan	Drip	Alfalfa	Drip	Pecan
-------	------	---------	------	-------



40-inch rows, 24-inch spacing

One plot, 20 feet; alleys 4 feet


Drip 8 inches both sides

A wide-angle photograph of a large agricultural field. The foreground and middle ground are filled with rows of young alfalfa seedlings, which appear as small green plants in the soil. The soil is dark brown and somewhat uneven. In the background, there is a long, low farm building with a dark roof. To the left of the building, there is a covered area, possibly a loading dock or a small shelter. Several vehicles, including white vans or trucks, are parked near the building. The sky is overcast and grey. The overall scene depicts a well-maintained agricultural field after transplanting.

Field after transplanting;
note seedling alfalfa,
2013

Cotton root rot (5-100%) in
6-month old alfalfa
20Nov2013

8 replications + borders

















U.S. Customs and
Border Protection
Office of Air and Marine
Uvalde Air Unit



Customs and
Protection
of Air and Marine
Air Unit



POLIOA

SILAF

Managing alfalfa







More CRR losses on west end; less disease in background

Data collection

- CRR in alfalfa (2013-2014 only)
- Dead seedlings – P.o. almost always on roots
- Height, lower stem diameter
- Bud break
- Survivors dug and examined

Data collection: disease

- Note dead seedlings
- Recover the roots
- Examine with magnification
- Record *Phymatotrichopsis omnivora*, root surface area (0, 0.01, 0.1, 1, $\geq 10\%$)

Recovery of roots from dead trees

- 2013,14,15 Hand digging
- 2016 Hand digging, air spade
- 2017 Excavator

**As seedlings
grew.....**





AIRMAN

AIRMAN
PD51855 T4

MMD
EQUIPMENT

Baker
Rentals
OVALDE, TEXAS
830-278-2894



Air-Spade, 90 p.s.i., Mach 2 air-jet





Root and column of soil exposed with Air Spade



Recover and bag dead roots from soil column










Statistics

	Dead-All		Dead-Fewer		Survivor-All		Survivor-Fewer	
	Entry	Prov	Entry	Prov	Entry	Prov	Entry	Prov
CRR	**	NS	**	NS	NS	NS	NS	NS
<i>P.o.</i> root area	NS	NS	NS	NS	NS	NS	NS	NS
Mortality	**	NS	**	NS	-	-	-	-


Dead plants with CRR, %

CRR 4/2017 P=0.009



Dead plants with CRR, %


CRR 4/2017 P=0.006



Entries with ≥ 65 plants


Dead plants with CRR, %

CRR 4/2017 P=0.30 N.S.




Dead plants with CRR, %

CRR 4/2017 P=0.26 N.S.




All dead plants, %

Mortality 4/2017 P=0.006




Mortality 4/2017 P=0.004



Entries with ≥ 65 plants


Diameter, mm


January 2017





Height, cm

January 2017









<http://cgru.usda.gov/carya/Manual/BUDBRK.html>

Bud break 1 to 9 index



Bud break 1 to 9 index





'Riverside' BWV 3-16 4 Apr 2017



87MX4-5.5 is a seedling grown from seed collected 8 Sept 1987 from this tree, growing in Ixmiquilpan, Hidalgo, Mexico, on the property of Norodino Rodriguiz. The original tree was 60 ft in height, 84 cm diameter, with moderate branching, no observed nutritional deficiencies, good health, heavy crop, when collected by Grauke, Payne, Wood and Parfitt.

Seedlings were planted in provenance orchards in College Station, TX and Byron, GA.

The Ixmiquilpan population is characterized by low, wide spreading canopies due to almost indeterminate terminal growth that begins early in the year and extends until very late, sometimes December.



87MX4-5.5, CSV 16-8

Seed from these two trees were planted in the Cotton Root Rot Nursery at Uvalde, TX.

87MX4-5.5 produces the largest seedlings in the test, but trees from Mexico (Southern Provenance) produce larger seedlings than US seedstocks.

Genomic tests are being run on 70 seedlings from each of these two mother trees to determine if they cross pollinated each other, and if pollen parent contributes to seedling vigor.

College Station provenance orchard 5 Apr 2017

87MX4-5.5

- Largest height and diameter
- Best in nematode screening
- Tolerates Na & Cl (Miyamoto)
- Performs well in inoculation with *Tuber lyonii*

GBS Sequencing

- Genotype By Sequence analysis
- Patricia Klein & Nolan Bentley
- 87MX5-1.7 pollen contributes to vigor of 87MX4-5.5 ?
- Establish relationship between genetic composition and field performance

Conclusions

- CRR screening is challenging, but possible
- No immunity in pecan, but some field resistance in seedling stage
- Rootstock choice should be considered for CRR management, and could become more important
- Consider tree size, fruitfulness, freeze risk, salt tolerance, water use, etc.